

Asthma + Allergies

June's focus is on asthma and allergies. This is the peak season for people with asthma and allergies and a great time to learn more about these conditions. Some people may have one or both of these conditions. According to the Asthma and Allergy Foundation of America,

- More than 25 million have asthma (Women are more likely to have asthma and die from it than men. It is more common in boys than girls.)
- Asthma is a leading chronic disease in children
- More than 50 million experience various types of allergies each year, and of these, about 25 million have seasonal allergic rhinitis (hay fever)
- Allergies are the sixth leading cause of chronic illness

Asthma and allergies are conditions that can occur at any age. The underlying cause is largely unknown, but both tend to run in families. Neither are curable for most patients, but both can be controlled with proper prevention and treatment. Discuss your symptoms with your Primary Care Provider—they are trained to manage most issues with asthma or allergies. They may refer you to an Allergist/Immunologist or Pulmonologist to help diagnosis or manage your condition. Whatever your medical team looks like, you should work together on your asthma or allergy treatment and management plan to help you feel and live better. An asthma control plan is one of the most important steps to managing your asthma—to keep you breathing well and avoid the ER and hospital care.

Participate in the following activities to learn more:

Activity #1:

Asthma vs. Allergies

Activity #2:

Spot the Differences

Activity #3:

Control Your Asthma

Activity #4:

Healthier Home Checklist



For questions or more information, email wellness@medica.com

Health Disparities

“Health disparities” are preventable differences in health among different social or racial groups. Health Disparities are a complex and challenging problem across the world. Social factors play the largest role in shaping the overall health of communities. For example, the ability to access healthcare, a safe living environment, racial equity, nutritious food, quality education, and supportive relationships free of discrimination are all incredibly important to your overall health.

People who face health disparities often have a higher risk for asthma and allergies. In the U.S., certain racial and ethnic groups experience a higher risk of asthma regardless of neighborhood income or population*.

Did You Know?

- Blacks and American Indian/Alaska Natives have the highest current asthma rates compared to other races and ethnicities. Blacks (10.9%) are 42 percent more likely than Whites (7.7%) to have asthma.**
- While asthma rates are relatively low for Hispanics overall, rates vary significantly among subgroups. Puerto Ricans in the continental United States have the highest current asthma rate of any racial or ethnic group.**
- African Americans have significantly higher adjusted odds of allergy to finfish.***

For more information, please visit:

- [*UW School of Medicine and Public Health](#)
- [**American Lung Association: Asthma Demographics](#)
- [***National Library of Medicine](#)

Activity 1: Asthma vs. Allergies

Test your knowledge. Read the statements below and guess which condition it applies to. Then learn more about these conditions on the educational chart on page 3. (Answers provided on page 4.)

	Asthma	Allergies
1. Most common condition affecting the human respiratory tract	<input type="checkbox"/>	<input type="checkbox"/>
2. Common treatments include epinephrine and/or antihistamines	<input type="checkbox"/>	<input type="checkbox"/>
3. Tobacco smoke is one of the most common triggers	<input type="checkbox"/>	<input type="checkbox"/>
4. Common symptoms: Wheezing, shortness of breath, and chest tightness	<input type="checkbox"/>	<input type="checkbox"/>
5. Occurs when a weak immune system responds to a foreign substance that is normally harmless	<input type="checkbox"/>	<input type="checkbox"/>
6. Common symptoms: Itchy/water eyes, sneezing, rashes, and wheezing	<input type="checkbox"/>	<input type="checkbox"/>
7. Leading cause of absence from school	<input type="checkbox"/>	<input type="checkbox"/>
8. Most common triggers are pollen, milk, and nuts	<input type="checkbox"/>	<input type="checkbox"/>
9. Can be controlled with regular use of anti-inflammatory medication	<input type="checkbox"/>	<input type="checkbox"/>

What’s the difference between asthma and allergies? Though they are often spoken of together, they are different conditions—allergies can commonly, but not always, provoke asthma symptoms. Learn more with this educational chart.

	Asthma	Allergies
Definition & Diagnosis	<p>A condition in which a person's airways become inflamed, become narrower due to swelling, and produce extra mucus, all of which makes it difficult to breathe.</p> <p>Diagnosed through physical exam and lung function tests (e.g., spirometry).</p>	<p>Abnormal and unusual response of the immune system to substances in the environment.</p> <p>Diagnosed through skin prick tests and blood tests.</p>
Organs Affected	Lungs and airways.	Multiple organs such as nose, lungs, throat, skin, lining of stomach, eyes and sinuses. Frequently allergies provoke asthma symptoms
Common Signs & Symptoms	<ul style="list-style-type: none"> • Shortness of breath • Coughing • Wheezing • Sensation of chest tightness 	<p>Symptoms vary widely and depend on how you're exposed. Some symptoms include:</p> <ul style="list-style-type: none"> • Itchy, watery eyes • Itchy and/or runny nose • Sneezing • Rashes • Hives (a rash with raised red patches) • Stomach cramps • Vomiting • Diarrhea • Bloating • Cough • Throat closing • Wheezing
Common Causes (can vary from person to person)	<p>Inflammation, mucus production and airway narrowing can start due to a variety of reasons:</p> <ul style="list-style-type: none"> • Viral infections • Allergies • Emotional upsets • Chemical irritants such as smoking or smoke exposure • Change in temperature or humidity • Exercise • Fatigue 	<p>Many different allergens are responsible for allergic reactions and include:</p> <ul style="list-style-type: none"> • Pollen • Dust • Food (e.g., fish, peanuts, shellfish, eggs) • Insect stings • Animal dander • Mold • Medications/Drugs • Latex



[Learn more](#) about Asthma Causes and Treatment.

[Learn more](#) about Allergy Prevention and Treatment.

[Learn more](#) about Allergy Test Types.

Activity 2: Spot the Differences

Can you spot the 10 differences between the asthma and allergen irritants in these two pictures? (*Answers provided on page 5.*)



Activity 4: Healthier Home Checklist

People spend more than 90% of their time indoors. Your Indoor air can be up to five times more polluted than outdoor air, per the U.S. Environmental Protection Agency (EPA), and an estimated 3.2 million deaths are attributed to household air pollution annually, according to the World Health Organization (WHO).

It is difficult to avoid problem-causing allergens entirely, but there are a few things you can do to lessen the impact they may have. Use this checklist for suggested tips on how to improve the indoor air quality of your home to lessen the impact of your allergies and/or asthma.

- Replace the air filters in your HVAC system as recommended by the manufacturer.
- Clean your living space once per week to remove dust, pollen, mold and animal dander.
- Select easy-to-clean home furnishings.
 - Choose blinds that can be wiped down over curtains. If you choose curtains, select fabric ones that can be washed in 130°F hot water regularly.
 - Vacuum furniture and mattresses weekly.
 - Use allergen-barrier mattress covers.
 - Wash throw blankets in 130° F hot water.
- Wash all bedding weekly in 130° F hot soapy water and dry on high heat. Replace pillows every 2 years. Use zippered allergen pillow protector covers.
- Leave your bed unmade to let it cool and air out. Warmth and moisture from your body encourage dust mite growth.
- Check your shower curtain for signs of mold/mildew and replace it, if necessary.
- Consider using a portable, room-sized air purifier.
- Keep windows closed during peak pollen season and when outdoor air pollution is high. Also, during peak pollen season, take your shoes off and change clothes upon coming in from the outdoors and take a shower before bed.
- For seasonal allergies, begin taking antihistamines in anticipation of your seasonal symptoms.
- As much as possible, avoid harsh chemicals like bleach, and remove items from your home with strong scents like candles.
- Measure and control the humidity inside your home. The Environmental Protection Agency (EPA) recommends that the indoor humidity range is between 30% to 50%.
- Improve airflow by leaving interior doors open and running exhaust fans in bathrooms and kitchens. Clean exhaust fan covers a few times per year.
- Consider using certified asthma and allergy friendly® products.

To learn more and to find certified asthma and allergy friendly® products check out the Asthma and Allergy Foundation of America.

Information obtained from SSM Health and The Asthma and Allergy Foundation of America (AAFA)