WellFirst Health

Adstiladrin® (nadofaragene firadenovec-vncg)

(Intravesical)

Document Number: IC-0691

Last Review Date: 02/02/2023 Date of Origin: 02/02/2023 Dates Reviewed: 02/2023

I. Length of Authorization

Coverage will be provided initially for 3 months and may be renewed every 6 months.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

- Adstiladrin suspension 3×10^{11} viral particles (vp)/mL- 20 mL single-dose vial: 4 vials every three months for four doses only

B. Max Units (per dose and over time) [HCPCS Unit]:

· 80 mL every 3 months x four doses

III. Initial Approval Criteria

Coverage is provided in the following conditions:

Patient is at least 18 years of age; AND

Universal Criteria¹

- · Patient does not have a hypersensitivity to interferon alfa; AND
- Patient is not immunosuppressed or immunodeficient; AND
- . Therapy will be used for intravesical instillation only; AND
- Must be used as a single agent; AND

Bladder Cancer † 1,4

- Patient has a diagnosis of non-muscle invasive bladder cancer (NMIBC) with carcinoma in situ *(with or without papillary tumors)*, **AND**
- Patient has high-risk disease that is unresponsive to Bacillus Calmette-Guerin (BCG) (defined as persistent disease following adequate BCG therapy [≥5 of 6 induction doses plus ≥2 doses of maintenance or of 2nd induction], disease recurrence after an initial tumor-free state following adequate BCG therapy, or T1 disease following a single induction course of BCG), AND



- Patient has undergone transurethral resection of bladder tumor (TURBT) to remove all resectable disease (Ta and T1 components); **AND**
- Patient does NOT have extra-vesical (i.e., urethra, ureter, or renal pelvis), muscle invasive (T2-T4), or metastatic urothelial carcinoma

FDA Approved Indication(s); C Compendia Recommended Indication(s); Φ Orphan Drug

IV. Renewal Criteria ^{1,4}

Coverage can be renewed based upon the following criteria:

- Patient continues to meet the universal and indication specific criteria as identified in section III; **AND**
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include the following: disseminated adenovirus infection, etc.; **AND**
 - <u>First Renewal</u>: Patient has a complete response (CR) to initial therapy (after 3 months) defined as a negative result for cystoscopy [with TURBT/biopsies as applicable] and urine cytology *: **OR**
 - o <u>Subsequent Renewals</u>: Patient has not experienced a high-grade or CIS recurrence

*Note: If patients with CIS do not have a complete response to treatment after 3 months or if CIS recurs, providers should consider cystectomy.

V. Dosage/Administration

Indication	Dose				
Bladder Cancer	The recommended dose of Adstiladrin is 75 mL at a concentration of 3 x 10 ¹¹ viral particles (vp)/mL by intravesical instillation once every three (3) months				
	• Premedication with an anticholinergic is recommended before each instillation.				
	*Note: Adstiladrin is not for intravenous use, topical use, or oral administration.				
	rin is a non-replicating adenoviral vector-based gene therapy. Follow universal biosafety ons for handling.				
	ndividuals who are immunosuppressed or immune-deficient, should not prepare, administer, or come nto contact with Adstiladrin.				
– Adstiladi	adrin is provided as a sterile frozen suspension.				
	y four (4) vials of Adstiladrin at room temperature (20°C to 25°C [68°F to 77°F]) until Adstiladrin is d. Do not expose the vials to higher temperatures. Protect from light.				
– Adstiladi	trin must be brought to room temperature (20°C to 25°C [68°F to 77°F]) prior to use.				
thawing	ne for thawing and bringing Adstiladrin to room temperature is approximately 8-10 h when g in the cardboard nest and approximately 3-5 h when thawing the vials outside the cardboard O NOT Refreeze.				
	the vial thawing procedure is initiated, the vials may be stored for up to 24 hours at room rature or refrigerated at 2°C to 8°C (36°F to 46°F). Visually inspect all 4 vials for visible particles				
	ADSTILADRIN [®] (nadofaragene firadenovec-vncg)				



and discoloration. The suspension is clear to slightly opalescent and may contain opalescent flecks. Do not use if visible particles or discoloration are observed. Mix gently. Do not shake.

VI. Billing Code/Availability Information

HCPCS Code:

J9999 - Not otherwise classified, antineoplastic drugs

NDC:

Adstiladrin suspension, with nominal concentration of 3 × 10¹¹ viral particles (vp)/mL in a carton of four frozen single-dose vials with an extractable volume of 20 mL/vial: 55566-1050-xx

VII. References

- 1. Adstiladrin [package insert]. Kuopio, Finland; Ferring Pharm, Inc; December 2022. Accessed December 2022.
- 2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for nadofaragene firadenovec. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed December 2022.
- 3. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium[®]) Bladder Cancer. Version 2.2022. National Comprehensive Cancer Network, 2022. The NCCN Compendium[®] is a derivative work of the NCCN Guidelines[®]. NATIONAL COMPREHENSIVE CANCER NETWORK[®], NCCN[®], and NCCN GUIDELINES[®] are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed December 2022.
- Boorjian SA, Alemozaffar M, Konety BR, et al. Intravesical nadofaragene firadenovec gene therapy for BCG-unresponsive non-muscle-invasive bladder cancer: a single-arm, openlabel, repeat-dose clinical trial. Lancet Oncol. 2021 Jan;22(1):107-117. doi: 10.1016/S1470-2045(20)30540-4. Epub 2020 Nov 27.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description	
C65.1	Malignant neoplasm of right renal pelvis	
C65.2	Malignant neoplasm of left renal pelvis	
C65.9	Malignant neoplasm of unspecified renal pelvis	

	ADSTILADRIN [®] (nadofaragene firadenovec-vncg)	
		Prior Auth Criteria
Page 3		Proprietary Information. Restricted Access – Do not disseminate or copy
		without approval.
		©2023, Magellan Rx Management



Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Articles (LCAs) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: https://www.cms.gov/medicare-coverage-database/search.aspx. Additional indications may be covered at the discretion of the health plan.

	Medicare Part B Administrative Contractor (MAC) Jurisdictions					
Jurisdiction	Applicable State/US Territory	Contractor				
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC				
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC				
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)				
6	MN, WI, IL	National Government Services, Inc. (NGS)				
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.				
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)				
N (9)	FL, PR, VI	First Coast Service Options, Inc.				
J (10)	TN, GA, AL	Palmetto GBA, LLC				
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC				
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.				
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)				
15	КҮ, ОН	CGS Administrators, LLC				

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

